



An Efficient Third Derivative Hybrid Block Method of Order Eleven for the Direct Solution of Second-Order Ordinary Differential Equation

D. Raymond^{1*}, Y. Skwame¹ and T. S. Kyagya²

¹Department of Mathematics and Statistics, Federal University, Wukari, Nigeria.

²Department of Mathematical Science, Adamawa State University, Mubi, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author DR designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors YS and TSK managed the analyses of the study. Author YS managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJOM/2018/40613

Editor(s):

(1) Nikolaos Dimitriou Bagis, Department of Informatics and Mathematics, Aristotelian University of Thessaloniki, Greece.

Reviewers:

(1) Grienggrai Rajchakit, Maejo University, Thailand.

(2) Suleyman Ogrekci, Amasya University, Turkey.

Complete Peer review History: <http://www.sciedomain.org/review-history/24212>

Received: 29th January 2018

Accepted: 12th April 2018

Published: 19th April 2018

Original Research Article

Abstract

This paper proposes an efficient third derivative hybrid block method of order eleven for solving second Order Ordinary Differential Equations directly. Method of interpolation and collocation of power series approximate solution to generate the continuous hybrid linear multistep method was used, which was evaluated at non-interpolating points to give a continuous block method. The discrete block method was recovered when the continuous block method was evaluated at all step points. The basic properties of the method were investigated and were found to be zero-stable, consistent, and convergent. The efficiency of the method was tested on some stiff equations and was found to give better approximation than the existing method compared with our result.

Keywords: Three-step; hybrid block method; third derivative; collocation; and interpolation method.

*Corresponding author: E-mail: raymond@fuwukari.edu.ng;

1 Introduction

This paper solves solution of stiff second order differential equation of the form

$$y'' = f(x, y(x), y'(x)), \quad y(x_0) = y_0, \quad y'(x_0) = y'_0 \quad (1)$$

Where f is continuous within the interval of integration. Different method have been proposed for the solution of (1) ranging from predictor-corrector method to hybrid methods. Despite the success recorded by the predictor-corrector methods, its major setback is that the predictor are in reducing order of accuracy especially when the value of the step-length is high and moreover the result are at overlapping interval. Direct methods of solving (1), which we shall employ has been discussed by many authors and they concluded that it is more convenient and accurate, among the authors that propose direct method are [1,2,3,4,5,6]. Block method have advantage of incorporating function evaluation at off-step points which afford the opportunity of circumventing the Dahlquist zero-stability barrier and it is actually possible to obtain convergent k-step methods of order $2k+1$. Hence, hybrid block method is less expensive in terms of number of function evaluation compare to predictor-corrector method, it also possess the properties of Runge-kutta for being self-starting and does not require starting values. Other author who proposes block methods are [7,8,9,10,11,12].

In this paper, we developed a three-step third derivative hybrid block method for direct solution of second Order Ordinary differential equations, which is implemented in block method. The method developed evaluates less function per step and circumventing the Dahlquist barrier's by introducing a hybrid points.

The paper is organised as follows: In section 2, we discuss the methods and the materials for the development of the method. Section 3 considers analysis of the basis properties of the method, which include convergence and stability region, numerical experiments where the efficiency of the derived method is tested on some stiff numerical examples and discussion of results. Lastly, we concluded in section 4.

2 Derivation of the Method

We consider a power series approximate solution of the form

$$y(x) = \sum_{j=0}^{2s+r-1} a_j \left(\frac{x-x_n}{h} \right)^j \quad (2)$$

where $r=2$ and $s=5$ are the numbers of interpolation and collocation points respectively, is considered to be a solution to (1).

The second and third derivative of (2) gives

$$\begin{aligned} y''(x) &= \sum_{j=2}^{2s+r-1} \frac{a_j j!}{h^2(j-2)} \left(\frac{x-x_n}{h} \right)^{j-2} \\ &= f(x, y, y'), \\ y'''(x) &= \sum_{j=3}^{2s+r-1} \frac{a_j j!}{h^3(j-3)} \left(\frac{x-x_n}{h} \right)^{j-3} \\ &= g(x, y, y'), \end{aligned} \quad (3)$$

Substituting (3) into (1) gives

$$f(x, y, y'') = \sum_{j=2}^{2s+r-1} \frac{a_j j!}{h^2(j-2)} \left(\frac{x-x_n}{h} \right)^{j-2} + \sum_{j=3}^{2s+r-1} \frac{a_j j!}{h^3(j-3)} \left(\frac{x-x_n}{h} \right)^{j-3} \quad (4)$$

Collocating (4) at all points $x_{n+s}, s = 0, \frac{1}{2}, 1, 2, 3$ and Interpolating Equation (2) at $x_{n+r}, r = 0, \frac{1}{2}$, gives a system of non linear equation of the form

$$AX = U \quad (5)$$

where

$$A = [a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, a_{11}]^T$$

$$U = \left[y_n, y_{\frac{n+1}{2}}, f_n, f_{\frac{n+1}{2}}, f_{n+1}, f_{n+2}, f_{n+3}, g_n, g_{\frac{n+1}{2}}, g_{n+1}, g_{n+2}, g_{n+3} \right]^T,$$

and

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & \frac{1}{2} & \frac{1}{4} & \frac{1}{8} & \frac{1}{16} & \frac{1}{32} & \frac{1}{64} & \frac{1}{128} & \frac{1}{256} & \frac{1}{512} & \frac{1}{1024} & \frac{1}{2048} \\ 0 & 0 & \frac{2}{h^2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{h^2} & \frac{3}{h^2} & \frac{3}{h^2} & \frac{5}{h^2} & \frac{15}{8h^2} & \frac{21}{16h^2} & \frac{7}{8h^2} & \frac{9}{16h^2} & \frac{45}{128h^2} & \frac{55}{256h^2} \\ 0 & 0 & \frac{2}{h^2} & \frac{6}{h^2} & \frac{12}{h^2} & \frac{20}{h^2} & \frac{30}{h^2} & \frac{42}{h^2} & \frac{56}{h^2} & \frac{72}{h^2} & \frac{90}{h^2} & \frac{110}{h^2} \\ 0 & 0 & \frac{2}{h^2} & \frac{12}{h^2} & \frac{24}{h^2} & \frac{40}{h^2} & \frac{60}{h^2} & \frac{84}{h^2} & \frac{112}{h^2} & \frac{144}{h^2} & \frac{180}{h^2} & \frac{220}{h^2} \\ 0 & 0 & \frac{2}{h^2} & \frac{18}{h^2} & \frac{36}{h^2} & \frac{60}{h^2} & \frac{90}{h^2} & \frac{126}{h^2} & \frac{168}{h^2} & \frac{216}{h^2} & \frac{270}{h^2} & \frac{330}{h^2} \\ 0 & 0 & 0 & \frac{6}{h^3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{6}{h^3} & \frac{12}{h^3} & \frac{15}{h^3} & \frac{15}{h^3} & \frac{105}{8h^3} & \frac{21}{2h^3} & \frac{63}{8h^3} & \frac{45}{8h^3} & \frac{495}{128h^3} \\ 0 & 0 & 0 & \frac{6}{h^3} & \frac{24}{h^3} & \frac{60}{h^3} & \frac{120}{h^3} & \frac{210}{h^3} & \frac{336}{h^3} & \frac{504}{h^3} & \frac{720}{h^3} & \frac{990}{h^3} \\ 0 & 0 & 0 & \frac{6}{h^3} & \frac{48}{h^3} & \frac{120}{h^3} & \frac{240}{h^3} & \frac{420}{h^3} & \frac{672}{h^3} & \frac{1008}{h^3} & \frac{1440}{h^3} & \frac{1980}{h^3} \\ 0 & 0 & 0 & \frac{6}{h^3} & \frac{72}{h^3} & \frac{180}{h^3} & \frac{360}{h^3} & \frac{630}{h^3} & \frac{1008}{h^3} & \frac{1512}{h^3} & \frac{2160}{h^3} & \frac{2970}{h^3} \end{bmatrix}$$

Solving (5) for a_i 's using Gaussian elimination method, gives a continuous hybrid linear multistep method of the form

$$P(x) = \sum_{j=0, \frac{1}{2}}^1 \alpha_j y_{n+j} + h^2 \left[\sum_{j=\frac{1}{2}}^1 \beta_j f_{n+j} + \sum_{j=0}^3 \beta_i f_{n+i} \right] + h^3 \left[\sum_{j=\frac{1}{2}}^1 \gamma_j g_{n+j} + \sum_{i=0}^3 \gamma_i g_{n+j} \right] \quad (6)$$

The coefficient of $y_{n+j}, j = 0, \frac{1}{2}$ and $f_{n+j}, j = 0, \frac{1}{2}, 1, 2, 3$ gives

where

$$\begin{aligned} a_0 &= \left(1 - \frac{2(x - x_n)}{h} \right), \quad a_{\frac{1}{2}} = \left(= \frac{2(x - x_n)}{h} \right) \\ \beta_0 &= \frac{23}{2970} \frac{(x - x_n)^{11}}{h^9} - \frac{148}{1215} \frac{(x - x_n)^{10}}{h^8} + \frac{6307}{7776} \frac{(x - x_n)^9}{h^7} - \frac{18017}{6048} \frac{(x - x_n)^8}{h^6} + \frac{14939}{2268} \frac{(x - x_n)^7}{h^5} - \frac{14497}{1620} \frac{(x - x_n)^6}{h^4} \\ &\quad + \frac{15563}{2160} \frac{(x - x_n)^5}{h^3} - \frac{139}{48} \frac{(x - x_n)^4}{h^2} + \frac{1}{2} (x - x_n)^2 - \frac{21654613}{153280512} (x - x_n)h \\ \beta_{\frac{1}{2}} &= \frac{4096}{185625} \frac{(x - x_n)^{11}}{h^9} - \frac{9856}{30375} \frac{(x - x_n)^{10}}{h^8} + \frac{59776}{30375} \frac{(x - x_n)^9}{h^7} - \frac{149312}{23625} \frac{(x - x_n)^8}{h^6} + \frac{161792}{14175} \frac{(x - x_n)^7}{h^5} - \frac{565376}{50625} \frac{(x - x_n)^6}{h^4} \\ &\quad + \frac{27392}{5625} \frac{(x - x_n)^5}{h^3} - \frac{256}{1125} \frac{(x - x_n)^4}{h^2} - \frac{2944883}{46777500} (x - x_n)h \\ \beta_1 &= -\frac{3}{110} \frac{(x - x_n)^{11}}{h^9} + \frac{37}{90} \frac{(x - x_n)^{10}}{h^8} - \frac{743}{288} \frac{(x - x_n)^9}{h^7} + \frac{975}{112} \frac{(x - x_n)^8}{h^6} - \frac{2855}{168} \frac{(x - x_n)^7}{h^5} + \frac{287}{15} \frac{(x - x_n)^6}{h^4} \\ &\quad - \frac{231}{20} \frac{(x - x_n)^5}{h^3} + 3 \frac{(x - x_n)^4}{h^2} - \frac{1255169}{28385280} (x - x_n)h \\ \beta_2 &= -\frac{7}{2970} \frac{(x - x_n)^{11}}{h^9} + \frac{8}{243} \frac{(x - x_n)^{10}}{h^8} - \frac{1459}{7776} \frac{(x - x_n)^9}{h^7} + \frac{3401}{6048} \frac{(x - x_n)^8}{h^6} - \frac{2171}{2268} \frac{(x - x_n)^7}{h^5} + \frac{1519}{1620} \frac{(x - x_n)^6}{h^4} \\ &\quad - \frac{361}{720} \frac{(x - x_n)^5}{h^3} + \frac{17}{144} \frac{(x - x_n)^4}{h^2} - \frac{1139959}{766402560} (x - x_n)h \\ \beta_3 &= -\frac{67}{371250} \frac{(x - x_n)^{11}}{h^9} + \frac{137}{60750} \frac{(x - x_n)^{10}}{h^8} - \frac{11207}{972000} \frac{(x - x_n)^9}{h^7} + \frac{11867}{378000} \frac{(x - x_n)^8}{h^6} - \frac{5611}{113400} \frac{(x - x_n)^7}{h^5} \\ &\quad + \frac{4627}{101250} \frac{(x - x_n)^6}{h^4} - \frac{1579}{67500} \frac{(x - x_n)^5}{h^3} + \frac{2}{375} \frac{(x - x_n)^4}{h^2} - \frac{6136241}{95800320000} (x - x_n)h \\ \gamma_0 &= \frac{1}{990} \frac{(x - x_n)^{11}}{h^8} - \frac{13}{810} \frac{(x - x_n)^{10}}{h^7} + \frac{281}{2592} \frac{(x - x_n)^9}{h^6} - \frac{205}{504} \frac{(x - x_n)^8}{h^5} + \frac{703}{756} \frac{(x - x_n)^7}{h^4} - \frac{361}{270} \frac{(x - x_n)^6}{h^3} \\ &\quad + \frac{173}{144} \frac{(x - x_n)^5}{h^2} - \frac{23}{36} \frac{(x - x_n)^4}{h} + \frac{1}{6} (x - x_n)^3 - \frac{1751221}{255467520} (x - x_n)h^2 \\ \gamma_{\frac{1}{2}} &= \frac{128}{12375} \frac{(x - x_n)^{11}}{h^8} - \frac{64}{405} \frac{(x - x_n)^{10}}{h^7} + \frac{2048}{2025} \frac{(x - x_n)^9}{h^6} - \frac{5536}{1575} \frac{(x - x_n)^8}{h^5} + \frac{6784}{945} \frac{(x - x_n)^7}{h^4} - \frac{29248}{3375} \frac{(x - x_n)^6}{h^3} \\ &\quad + \frac{2176}{375} \frac{(x - x_n)^5}{h^2} - \frac{128}{75} \frac{(x - x_n)^4}{h} + \frac{51643}{1559250} (x - x_n)h^2 \\ \gamma_1 &= \frac{1}{110} \frac{(x - x_n)^{11}}{h^8} - \frac{2}{15} \frac{(x - x_n)^{10}}{h^7} + \frac{233}{288} \frac{(x - x_n)^9}{h^6} - \frac{587}{224} \frac{(x - x_n)^8}{h^5} + \frac{39}{8} \frac{(x - x_n)^7}{h^4} - \frac{125}{24} \frac{(x - x_n)^6}{h^3} + 3 \frac{(x - x_n)^5}{h^2} \\ &\quad - \frac{3}{4} \frac{(x - x_n)^4}{h} + \frac{59057}{5677056} (x - x_n)h^2 \end{aligned}$$

$$\begin{aligned}\gamma_2 &= \frac{1}{990} \frac{(x-x_n)^{11}}{h^8} - \frac{11}{810} \frac{(x-x_n)^{10}}{h^7} + \frac{193}{2592} \frac{(x-x_n)^9}{h^6} - \frac{31}{144} \frac{(x-x_n)^8}{h^5} + \frac{269}{756} \frac{(x-x_n)^7}{h^4} - \frac{46}{135} \frac{(x-x_n)^6}{h^3} \\ &\quad + \frac{43}{240} \frac{(x-x_n)^5}{h^2} - \frac{1}{24} \frac{(x-x_n)^4}{h} + \frac{131671}{255467520} (x-x_n) h^2 \\ \gamma_3 &= \frac{1}{24750} \frac{(x-x_n)^{11}}{h^8} - \frac{1}{2025} \frac{(x-x_n)^{10}}{h^7} + \frac{161}{64800} \frac{(x-x_n)^9}{h^6} - \frac{337}{50400} \frac{(x-x_n)^8}{h^5} + \frac{79}{7560} \frac{(x-x_n)^7}{h^4} - \frac{259}{27000} \frac{(x-x_n)^6}{h^3} \\ &\quad + \frac{11}{2250} \frac{(x-x_n)^5}{h^2} - \frac{1}{900} \frac{(x-x_n)^4}{h} + \frac{12119}{912384000} (x-x_n) h^2\end{aligned}$$

Where $t = \frac{x-x_n}{h}$, $y_{n+j} = y(x_n + jh)$ and $f_{n+j} = f((x_n + jh), y(x_n + jh), y'(x_n + jh))$

Differentiating (6) once yields

$$p'(x) = \frac{1}{h} \sum_{j=0, \frac{1}{2}} \alpha_j y_{n+j} + h \left[\sum_{j=\frac{1}{2}} \beta_j f_{n+j} + \sum_{j=0}^3 \beta_j f_{n+j} \right] + h^2 \left[\sum_{j=\frac{1}{2}} \gamma_j g_{n+j} + \sum_{j=0}^3 \gamma_j g_{n+j} \right] \quad (7)$$

Equation (6) is evaluated at non-interpolating points, and evaluating (7) at all points gives a discrete block formula of the form

$$A^{(0)} Y_m^{(i)} = \sum_{i=0}^1 h^i e_i y_n^{(i)} + h^2 b_i f(y_n) + h^2 d_i f(Y_m) + h^3 c_i g(y_n) + h^3 r_i g(Y_m) \quad (8)$$

Where

$$\begin{aligned}Y_m &= \left[y_{\frac{n+1}{2}}, y_{n+1}, y_{n+2}, y_{n+3} \right]^T, \quad f(y_m) = \left[f_{\frac{n+1}{2}}, f_{n+1}, f_{n+2}, f_{n+3} \right]^T, \quad g(y_m) = \left[g_{\frac{n+1}{2}}, g_{n+1}, g_{n+2}, g_{n+3} \right]^T \\ y_n^{(i)} &= \left[y_{n-1}^{(i)}, y_{n-2}^{(i)}, y_{n-3}^{(i)}, y_n^{(i)} \right]^T, \quad f(y_n) = \left[f_{n-1}, f_{n-2}, f_{n-3}, f_n \right]^T, \quad g(y_n) = \left[g_{n-1}, g_{n-2}, g_{n-3}, g_n \right]^T\end{aligned}$$

And $A^{(0)} = 8 \times 8$ identity matrix.

when $i = 0$

$$\begin{aligned}e_0 &= \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad e_i = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 3 \end{bmatrix}, \quad b_0 = \begin{bmatrix} 0 & 0 & 0 & \frac{21654613}{306561024} \\ 0 & 0 & 0 & \frac{41257}{249480} \\ 0 & 0 & 0 & \frac{5399}{13365} \\ 0 & 0 & 0 & \frac{1233}{1232} \end{bmatrix}, \quad d_0 = \begin{bmatrix} \frac{2944883}{93555000} & \frac{1255169}{56770560} & \frac{1139959}{1532805120} & \frac{6136241}{1916006400} \\ \frac{36416}{155925} & \frac{243}{2464} & \frac{589}{249480} & \frac{493}{4989600} \\ \frac{1650688}{1670625} & \frac{248}{495} & \frac{1411}{13365} & \frac{2312}{1670625} \\ \frac{159552}{7047} & \frac{7047}{4167} & \frac{4167}{122211} & \frac{122211}{1540000} \end{bmatrix} \\ c_0 &= \begin{bmatrix} 0 & 0 & 0 & \frac{1751221}{510935040} \\ 0 & 0 & 0 & \frac{137}{15840} \\ 0 & 0 & 0 & \frac{778}{31185} \\ 0 & 0 & 0 & \frac{1053}{12320} \end{bmatrix}, \quad r_0 = \begin{bmatrix} -\frac{51643}{3118500} & -\frac{59057}{11354112} & -\frac{131671}{510935040} & -\frac{12119}{1824768000} \\ -\frac{763}{17325} & -\frac{529}{27720} & -\frac{269}{332640} & -\frac{17}{831600} \\ \frac{2048}{779625} & \frac{404}{3465} & \frac{652}{31185} & \frac{212}{779625} \\ \frac{864}{1215} & \frac{1215}{1647} & \frac{1647}{801} & -\frac{801}{77000} \end{bmatrix}\end{aligned}$$

when $i = 1$

$$e_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}, b_0 = \begin{bmatrix} 0 & 0 & 0 & \frac{12976613}{69672960} \\ 0 & 0 & 0 & \frac{52363}{272160} \\ 0 & 0 & 0 & \frac{2497}{8505} \\ 0 & 0 & 0 & \frac{1161}{1120} \end{bmatrix}, d_0 = \begin{bmatrix} \frac{245039}{1063125} & \frac{207791}{2580480} & \frac{182827}{69672960} & \frac{975887}{8709120000} \\ \frac{506752}{45056} & \frac{659}{76} & \frac{1061}{685} & \frac{5311}{124} \\ \frac{42525}{18048} & \frac{315}{3321} & \frac{1701}{519} & \frac{42525}{47589} \\ \frac{4375}{1120} & \frac{1120}{1120} & \frac{1120}{1120} & \frac{140000}{140000} \end{bmatrix}$$

$$c_0 = \begin{bmatrix} 0 & 0 & 0 & \frac{234319}{23224320} \\ 0 & 0 & 0 & \frac{977}{90720} \\ 0 & 0 & 0 & \frac{13}{567} \\ 0 & 0 & 0 & \frac{129}{1120} \end{bmatrix}, r_0 = \begin{bmatrix} -\frac{37879}{567000} & -\frac{139}{3360} & -\frac{3007}{3317760} & -\frac{13481}{580608000} \\ -\frac{2944}{70875} & -\frac{139}{3360} & -\frac{17}{12960} & -\frac{73}{2268000} \\ -\frac{2048}{14175} & -\frac{32}{105} & -\frac{23}{405} & -\frac{8}{14175} \\ -\frac{1152}{875} & -\frac{1863}{1120} & -\frac{63}{160} & -\frac{921}{28000} \end{bmatrix}$$

3 Analysis of Basic Properties of the Method

3.1 Order of the block

Let the linear operator $L\{y(x); h\}$ associated with the discrete block method (8) be defined

$$L\{y(x); h\} = A^{(0)}Y_m^i - \sum_{j=0}^3 h^i e_j y_n^i - h^2(b_0 f(y_n) + d_0 F(Y_m)) - h^3(c_0 g(y_n) + G(Y_m)) \quad (9)$$

Expanding (9) in Taylor series and comparing the coefficient of h gives

$$L\{y(x); h\} = C_0 y(x) + C_1 y'(x) + \dots + C_p h^p y^p(x) + C_{p+1} h^{p+1} y^{p+1}(x) + C_{p+2} h^{p+2} y^{p+2}(x) + \dots$$

Definition: Linear operator L and associated block formula are said to be of order

p , if $C_0 = C_1 = \dots = C_p = C_{p+1} = 0$, and $C_{p+2} \neq 0$. C_{p+2} called the error constant and implies that the truncation error is given by $t_{n+k} = C_{p+2} h^{p+2} y^{p+2}(x) + O(h^{p+3})$

For our method, expanding (8) in Taylor series

and comparing the coefficient of h gives $C_0 = C_1 = C_2 = C_3 = \dots = C_{11} = 0$ and

$$C_{12} = \left[\frac{238807}{206009008128000}, \frac{233}{67060224000}, \frac{13}{449064000}, \frac{87}{275968000}, \frac{51643}{12875563008000}, \frac{529}{100590336000}, \frac{163}{3143448000}, \frac{89}{137984000} \right]^T$$

3.2 Zero stability of our method

Definition: A block method is said to be zero-stable if as $h \rightarrow 0$, the root $z_i, i=1(k)$ of the first characteristic polynomial $\rho(z)=0$ that is $\rho(z) = \det \left[\sum_{j=0}^k A^{(i)} z^{k-i} \right] = 0$ Satisfies $|z_i| \leq 1$ and for those roots with $|z_i|=1$, multiplicity must not exceed two. The block method for $k=3$, with one off grid collocation point expressed in the form

$$\rho(z) = z \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & \frac{h}{2} \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & h \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 2h \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 3h \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} = z^6(z-1)^2$$

$$\rho(z) = z^6(z-1)^2 = 0,$$


Hence, our method is zero-stable.

3.3 Region of absolute stability of the two step one offstep point

We shall adopt the boundary locus method to determine the region of absolute stability of the implicit three-step third derivative hybrid block method. This gives stability polynomial below

$$\begin{aligned} \bar{h}(w) = & h^{12} \left(\left(\frac{8581}{299376000000} \right) w^4 - \left(\frac{23329}{34927200000} \right) w^3 \right) + h^{11} \left(\left(\frac{969161}{4191264000000} \right) w^4 + \left(\frac{25354921}{1047816000000} \right) w^3 \right) \\ & - h^{10} \left(\left(\frac{13481}{93139200000} \right) w^4 + \left(\frac{349245913}{838252800000} \right) w^3 \right) + h^9 \left(\left(\frac{20731903}{1077753600000} \right) w^4 + \left(\frac{172960967}{1077753600000} \right) w^3 \right) \\ & - h^8 \left(\left(\frac{177004147}{1077753600000} \right) w^4 + \left(\frac{22083914003}{1077753600000} \right) w^3 \right) + h^7 \left(\left(\frac{464273}{907200000} \right) w^4 - \left(\frac{653723467}{39916800000} \right) w^3 \right) \\ & + h^6 \left(\left(\frac{2144917}{1995840000} \right) w^4 + \left(\frac{306556939}{1995840000} \right) w^3 \right) - h^5 \left(\left(\frac{19800841}{1496880000} \right) w^4 + \left(\frac{420829301}{598752000} \right) w^3 \right) \\ & + h^4 \left(\left(\frac{4569569}{166320000} \right) w^4 - \left(\frac{74560469}{166320000} \right) w^3 \right) + h^3 \left(\left(\frac{7457}{111375} \right) w^4 - \left(\frac{673693}{356400} \right) w^3 \right) \\ & - h^2 \left(\left(\frac{561383}{1782000} \right) w^4 + \left(\frac{7457617}{1782000} \right) w^3 \right) + w^4 - w^3 \end{aligned}$$

The absolute stability region of the new method is plotted and shown below



3.4 Numerical example

Problem I. We consider a highly stiff problem

$$y'' + 1001y' + 1000y, \quad y(0) = 1, y'(0) = -1$$

Exact Solution: $y(x) = \exp(-x)$, $h = \frac{1}{10}$

x-values	Exact solution	Computed solution	Error in our method	Error in [8]
0.100	0.90483741803595957316	0.90483741803595957382	6.600000E(-19)	1.054712E(-14)
0.200	0.81873075307798185867	0.81873075307798185402	4.650000E(-18)	1.776357E(-14)
0.300	0.74081822068171786607	0.74081822068171786407	2.000000E(-18)	2.342571E(-14)
0.400	0.67032004603563930074	0.67032004603563929888	1.860000E(-18)	2.797762E(-14)
0.500	0.60653065971263342360	0.60653065971263342181	1.790000E(-18)	3.130829E(-14)
0.600	0.54881163609402643263	0.54881163609402643074	1.890000E(-18)	3.397282E(-14)
0.700	0.49658530379140951470	0.49658530379140951247	2.230000E(-18)	3.563816E(-14)
0.800	0.44932896411722159143	0.44932896411722158881	2.620000E(-18)	3.674838E(-14)
0.900	0.40656965974059911188	0.40656965974059910872	3.160000E(-18)	3.730349E(-14)
1.00	0.36787944117144232160	0.36787944117144231770	3.900000E(-18)	3.741452E(-14)

Problem II. $f(x, y, y') = y'$, $y(0) = 0$, $y'(0) = -1$.

Exact Solution: $y(x) = 1 - e^x$ with, $h = \frac{1}{100}$

x-values	Exact solution	Computed solution	Error in our method	Error in [10]
0.01	-0.1051709180756476248	-0.10517091807564762481	1.00000E(-20)	8.32667E(-17)
0.02	-0.2214027581601698339	-0.22140275816016983388	2.00000E(-20)	2.775558(-16)
0.03	-0.3498588075760031040	-0.34985880757600310361	3.90000E(-19)	5.551116(-16)
0.04	-0.4918246976412703178	-0.49182469764127031664	1.16000E(-18)	9.436896(-16)
0.05	-0.6487212707001281468	-0.64872127070012814473	2.07000E(-18)	2.109425(-15)
0.06	-0.8221188003905089749	-0.82211880039050897131	3.59000E(-18)	3.219647(-15)
0.07	-1.0137527074704765216	-1.01375270747047651580	5.80000E(-18)	4.440892(-15)
0.08	-1.2255409284924676046	-1.22554092849246759630	8.30000E(-18)	5.995204(-15)
0.09	-1.4596031111569496638	-1.45960311115694965230	1.15000E(-17)	7.771561(-15)
0.10	-1.71828182845904523540	-1.71828182845904521940	1.60000E(-17)	1.065814(-14)

4 Conclusions

It is evident from the above tables that our proposed methods is indeed accurate, and can handle stiff equations. Also in terms of stability analysis, the method is *A-stable*.

Comparing the new method with the existing method [8,10],the result presented in the tables 1 and 2 shows that the new method performs better than the existing method [8,10]. In this article, a three-step block method with one off-step point is derived via the interpolation and collocation approach. The developed method is consistent, convergent and zero-stable with an A-Stable region and order eleven.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Awoyemi DO. A class of hybrid collocation method for third order ordinary differential equations. Intern. J. Compu. Math. 2007;82(10):1287–1293.
DOI: 10.1080/00207160500112902
- [2] Adesanya AO, Anake TA, Udoh MO. Improved continuous method for direct solution of general second order ordinary differential equation. J. of Nig. Assoc. of Maths. Phys. 2008;3:59-62.
- [3] Kayode SJ, Adeyeye AO. A 3-step hybrid method for the direct solution of second order initial value problem. Australian J. of Basic and Applied Sciences. 2001;5(12):2121-2126.
- [4] Awoyemi DO, Adebile EA, Adesanya OA, Anake TA. Modified block for the direct of second order ordinary differential equations. Intern. J. Applied Maths & Comp. 2011;3(3):181–188.
- [5] Adeniran AO, Ogundare BS. An efficient hybrid numerical scheme for solving general second order initial value problems (IVPs). International Journal of Applied Mathematical Research. 2015;4(2):411-419.
- [6] Kayode SJ. A zero-stable method for direct solution of fourth order ordinary differential equation. American J. of Applied Sciences. 2009;5(11):1461-1466.
- [7] Abdelrahim R, Zurni O. Direct solution of second-order ordinary differential equation using a single-step hybrid block method of order five. Mathematical and Computational Applications. 2016;21:12.

- [8] Mohammad A, Zurni O. Implicit one-step block hybrid third-derivative method for the direct solution of initial value problems of second-order ordinary differential equations. Hindawi Journal of Applied Mathematics; 2017. Article ID: 8510948.
Available:<https://doi.org/10.1155/2017/8510948>
- [9] Adesanya AO, Alkali MA, Joshua S. Order five hybrid block method for the solution of second order ordinary differential equation. 2014;8(3):285-295.
- [10] Mohammad A, Zurni O. Generalized two-hybrid one-step implicit third derivatives block method for the direct solution of second order ordinary differential equations. International Journal of Pure and Applied Mathematics. 2017;112(3):497-517.
- [11] Anake TA, Awoyeni DO, Adesanya AO. A one step method for the solution of general second order ordinary differential equations. Intern. J. Sci. and Engin. 2012;2(4):159-163.
- [12] Fatunla SO. Block methods for second order IVPs. Int. J. Comput. Maths. 1991;41:55-63.

© 2018 Raymond et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)
<http://www.sciedomain.org/review-history/24212>