

Growth and Quality of Papaya ‘Golden THB’ Seedlings Cultivated under Different Irrigation Depths

**Robson Prucoli Posse^{1*}, Francielly Valani¹, Vinicius de Souza Oliveira²,
Leandro Moreira da Silva¹, Stefany Sampaio Silveira¹,
Sophia Machado Ferreira da Silva¹ and Geilson Silva Costa¹**

¹Federal Institute of Espírito Santo – Campus Itapina, Highway BR-259, Km 70, Countryside, Post Office Box 256, CEP: 29709-910, Colatina, Espírito Santo, Brazil.

²Federal University of Espírito Santo – Center North of Espírito Santo, Highway BR-101, Km 60, District Litorâneo, CEP: 29932-540, São Mateus, Espírito Santo, Brazil.

Authors' contributions

This work was carried out in collaboration between all authors. Author RPP designed the study, conducted the experiment in the field with authors FV, VSO, LMS, SSS and SMFS managed the writing of the manuscript. Authors RPP and FV managed the analyses of study. Authors RPP and VSO performed the statistical analysis. Authors GSC and RPP performed translation of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/46286

Editor(s):

(1) Dr. Abdel Razik Ahmed Zidan, Professor, Hydraulics and Water Resources, Mansoura University, Egypt.

Reviewers:

(1) Saleh Ahmed Shahriar, Sher-e-Bangla Agricultural University, Bangladesh.

(2) Charles Afriyie-Debrah, CSIR-Crops Research Institute, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46286>

Original Research Article

Received 10 October 2018

Accepted 25 December 2018

Published 15 January 2019

ABSTRACT

The obtainment of seedlings is one of the main stages of the papaya crop and good irrigation management provides proper water status, contributing to higher quality seedlings. Thus, this work had as objective to evaluate the growth and quality of Papaya 'Golden THB' seedlings submitted to different irrigation depths. The study was carried out at the Federal Institute of Espírito Santo, Campus Itapina, located in Colatina, Northwest region of the State of Espírito Santo, Brazil, during the period from May 16 to July 19, 2018. The experimental design was completely randomized with six treatments composed of 25 plants each, totaling 150 seedlings in the experiment. The treatments consisted in the application of six daily irrigation depths, corresponding to 4, 6, 8, 10, 12

*Corresponding author: E-mail: rpposse@gmail.com;

and 14 mm. At 65 days after planting, the seedlings were evaluated for morphological characteristics: length of the root system, leaf area, plant height, stem diameter, dry mass of the aerial part, dry mass of the root system, total dry mass and Dickson quality index. Irrigation depths interfered in the growth and quality of papaya 'Golden THB' seedlings during their production period. The recommended irrigation depth for the production of papaya 'Golden THB' seedlings with high quality is 8.15 mm d^{-1} , which showed a higher Dickson quality index and good responses in all the morphological characteristics studied.

Keywords: *Carica papaya L.*; efficiency of water use; irrigation management; Dickson index; vegetative development; water replenishment.

1. INTRODUCTION

The papaya (*Carica papaya L.*) is one of the most cultivated and consumed fruit trees in tropical and subtropical regions, and its production is widespread in almost all tropical America countries. The cultivar 'Golden THB' belongs to the 'Solo' group, being characterized by high productivity, uniformity and vigor in the plantations, being destined mainly for the export market [1].

One of the main stages to reach the maximum genetic potential of papaya growth is to obtain seedlings [2]. The correct management of irrigation of the seedlings maintains an adequate water status contributing to get plants with a higher quality standard [3]. In many cases, the application of water to the plants is done empirically, without technical knowledge, leading to a reduction of the productive potential of the crops [4].

The amount of water is a limiting factor in the development of plants and lack or excess affects plant health, growth and production [5]. The lack of water creates water stress and a decrease in the absorption of nutrients. The excess is conducive to disease development, causes leaching of nutrients and excessive water consumption, causing socioenvironmental problems [6]. So, the knowledge of the set of techniques for the production of seedlings favors the elaboration of more efficient protocols for making seedlings with higher quality [7].

Thus, studies are necessary to evaluate the efficiency of water use in the development of papaya seedlings. The objective of this work was to evaluate the growth and quality of papaya 'Golden THB' seedlings submitted to different irrigation depths.

2. MATERIALS AND METHODS

The study took place at the Federal Institute of Espírito Santo, Campus Itapina, located in

Colatina, Northwest region of the State of Espírito Santo, Brazil, located at $19^{\circ}29'$ South, $40^{\circ}45'$ West and altitude of 62 meters, from May 16 to July 19, 2018. The climate of the region, according to Köppen's classification, is Tropical Aw [8]. The zone is characterized by the irregularity of the rain and the occurrence of high temperatures.

The experiment was conducted in an agricultural greenhouse covered with transparent plastic film and black polypropylene screen with 50% shading, linear dimensions of 25 m x 5 m and height of 3 m, in six individual environments (BOX) with dimensions of 2.20 m long by 1.10 m wide each, isolated by transparent plastic canvas on the sides. Each BOX consisted of six GREEN MIST anti-mist nebulizers from NaanDanJain[®], located 1 m above the seedlings and spaced 0.8 m apart, with a daily watering frequency of 10 hours (from 07:00 am 05:00 pm), operating at a service pressure of 2 kgf cm^{-2} , individually controlled by electronic controllers and centrifugal pumps of 0.5 hp independently installed.

The treatments consisted in the application of six daily irrigation depths, corresponding to 4, 6, 8, 10, 12 and 14 mm d^{-1} , for the production of papaya seedlings. The experimental design was completely randomized, with each treatment consisting of 25 plants, totaling 150 seedlings in the experiment.

For the production of the seedlings were used seeds of the papaya 'Golden THB' obtained from the company NorteFru[®] located in Linhares-ES. The seedlings were produced in tubes with a top diameter of 53 mm, 190 mm in height and 280 ml of volumetric capacity. The tubes were arranged in holders with 54 cells, distributed alternately for not limiting the light in the seedlings, which could lead to them to a stretching. All tubes were previously sanitized and sterilized with 2% sodium hypochlorite diluted in water and filled with Tropstrato HT[®] Vegetable substrate plus

Osmocote Plus® 15-9-12 (3M), at a dosage of 3g tube⁻¹, which had the following chemical composition: N = 15%, (7% ammonia and 8% nitrate), P₂O₅ = 9%, K₂O = 12%, Mg = 1.3%, S = 5.9%, Cu = 0.05%, Fe = 0.46%, Mn = 0.06% and Mo = 0.02%. No additional fertilization was done.

During the execution of the experiment, variations in temperature and relative humidity within the greenhouse were monitored by a WatchDog® Model 200 Data Logger. Externally climatic variables were recorded by an ONSET® weather station, reference evapotranspiration (ET_o) was estimated by the Penman-Monteith method FAO-56 Standard [9] by the equation:

$$ET_o = \frac{0.408\Delta(Rn - G) + \gamma \frac{900}{T + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34 u_2)}$$

in which ET_o is the daily reference evapotranspiration (mm.d⁻¹); Rn is the daily radiation balance (MJ.m⁻².d⁻¹); G is the daily flow of heat in the soil (MJ.m⁻².d⁻¹); T is the daily average air temperature (°C); u₂ is the daily average wind speed at 2 m in height (m.s⁻¹); e_s is the saturation pressure of the daily average water vapor (kPa); e_a is the daily average water vapor pressure (kPa); Δ is the slope of the vapor pressure curve at the point of T (kPa.°C⁻¹) and γ is the psychrometric coefficient (kPa.°C⁻¹).

At 65 days after planting, the seedlings were evaluated for the following morphological characteristics: a) leaf area (LA), expressed in cm², estimated with LI-COR leaf area measurer model LI-3100C; b) length of the root system (LRS), measured from the base of the stem to the largest root length, in cm, with graduated ruler; c) plant height (PH), measured in cm, with the use of a graduated ruler, from the stem to the apical bud; d) stem diameter (SD), measured 2 cm above the edge of the tube, in mm, with a digital caliper from Metrotools, model MPD-150; e) dry mass of the aerial part (DMAP), dry mass of the root system (DMRS) and total dry mass (TDM), expressed in grams, through an electronic scale with an accuracy of 0.001g; f) Dickson quality index (DQI) according to Dickson et al. [10], given by:

$$DQI = \frac{TDM}{\frac{PH}{SD} + \frac{DMAP}{DMRS}}$$

To obtain the DMAP and DMRS, the plants were subjected to drying process in greenhouse with forced circulation of air at 65°C for 72 hours and weighed on electronic scale with precision of 0.001 g. The TDM was obtained by the sum of DMAP and DMRS.

The data were subjected to analysis of variance by the F test, at 5% probability, when significant, regression models were adjusted to better explain the effect of the irrigation depths on the characteristics analyzed by the R software [11]. The maximum points were determined by the first derivative of the regression equations.

3. RESULTS AND DISCUSSION

During the experimental period, the external meteorological variations of the agricultural greenhouse presented a maximum temperature of 22.0 to 34.2°C (average of 29.78°C), average of 19.7 to 24.8°C (average of 22.18°C) and minimum of 13.1 to 20.6°C (average of 16.98°C) (Fig. 1). The average relative humidity varied from 76.9 to 94.1% (average of 82.78%) (Fig. 2). The maximum reference evapotranspiration observed was 2.97, minimum of 1.06 and average of 2.15 mm d⁻¹ (Fig. 3).

Within the agricultural greenhouse, the maximum temperature varied from 25.2 to 45.4°C (average of 38.84°C), the average temperature had variations from 20.9 to 27.5°C (average of 24.66°C) and the minimum temperature ranged from 14.1 to 21.3°C (average of 17.68°C) (Fig. 4). The average relative humidity ranged from 60.8 to 80.4% (average of 65.76%) (Fig. 5). These conditions of average temperature and relative humidity are considered ideal for the development of papaya plants [12]. On average, the maximum, average and minimum temperatures within the agricultural greenhouse were 30.42, 11.18 and 4.12% higher than the external averages, while relative humidity decreased by 20.6%.

After analysis of variance, a significant difference ($P < 0.05$) was observed for all the evaluated characteristics, noting that the application of the different irrigation depths interfered in the growth and quality of the papaya 'Golden THB' seedlings.

The length of the root system (LRS) showed a quadratic effect, with determination coefficient (R^2) of 0.745 and maximum point with 27.78 cm

in the irrigation depth of 9.44 mm d⁻¹ (Fig. 6A). Leaf area (LA) presented a quadratic adjustment, with its maximum point in the irrigation depth of 7.28 mm d⁻¹, with an area of 116.65 cm² and R² of 0.9401 (Fig. 6B). Measure LA is very important because the leaves are responsible for capturing

the highest amount of solar energy required for photosynthesis, producing photosimilates used for maintenance and growth of the plant [13,14]. So, higher AF indicates more suitable seedlings the field conditions because they possess larger photosynthetically active area [15].




Fig. 1. Maximum, minimum and average external temperatures during the experimental period in Colatina, Espírito Santo




Fig. 2. External average relative humidity during the experimental period in Colatina, Espírito Santo




Fig. 3. Reference evapotranspiration during the experimental period in Colatina, Espírito Santo




Fig. 4. Maximum, minimum and average temperatures within the agricultural greenhouse during the experimental period in the municipality of Colatina, Espírito Santo

Under stress conditions, plants change their morphological characteristics [16]. The water deficit caused by the lack of water in the smaller irrigation depths decreases the swelling pressure of the plant cells, reflecting in turgidity dependent processes, as the expansion of the leaf area and the extension of the root system. The expansion of the leaf area is early affected when the amount of water is reduced, this morphological change leads the plants to transpire less,

supporting for a longer time the availability of water in the substrate [17].

On the other hand, the excess of irrigation affected the length of the root system and the leaf area of the seedlings. Under such conditions plants are exposed to an oxygen-depleted environment (anoxia). Virtually all plants need oxygen to perform their metabolic activities, thus under anoxic conditions their development is

limited. The root respiration rate is diminished in anaerobic environments, causing damage to its growth. In addition, the roots cannot supply the energy demand required to sustain the physiological processes of the aerial part of the plants, limiting their development [17].

The plant height (PH) presented quadratic adjustment, with R^2 of 0.7697. The maximum height of 16.01 cm was observed in the irrigation depth of 8.84 mm d^{-1} (Fig. 7A). The lowest heights were observed in the largest and the shortest irrigation depth. According to Delgado et

al. [7], the excess water provides greater leaching of nutrients present in the substrate, limiting the size of the plants. On the other hand, lack of water prevents the expansion of the cell wall due to altered hormonal balance and cellular turgidity impairing plant growth [18].

The height is used as a base parameter for the transplanting of seedlings to the field, taking advantage of larger seedlings, since seedlings of smaller stature can be constantly damaged, impairing their development and production [5].




Fig. 5. Average relative humidity within the agricultural greenhouse during the experimental period in Colatina, Espírito Santo




Fig. 6. Length of root system (A) and leaf area (B) of papaya 'Golden THB' seedlings submitted to different irrigation depths

The stem diameter (SD) presented quadratic behavior, with a maximum point of 6.04 mm in the irrigation depth of 6.29 mm d⁻¹ and R² of 0.9225 (Fig. 7B). The behavior of SD was similar to that presented by LA. This fact can be explained, according to Melo et al. [15], because of the high predictive correlation, thus, plants with larger leaf area produce more photosimilates and, consequently, larger stem diameter, the opposite is observed in plants with smaller leaf area that has reduced stem diameter.

Stem diameter and plant height are very important characteristics, since they are related to possible tipping of seedlings in the field and they are easily evaluated, expressing quality in every stage of the seedling development [19].

The dry mass of the aerial part (DMAP), dry mass of the root system (DMRS) and total dry mass (TDM) showed a quadratic effect, with a maximum point of 0.95 g, 0.44 g and 1.39 g irrigation depths of 8.36, 8.52, 8.42 mm d⁻¹ and R² of 0.9607, 0.8607 and 0.9581, respectively (Fig. 8A, 8B and 8C). There is similarity in the adjustments of LA, DMAP, DMRS and TDM, so it can be observed that the increase in foliated area allowed greater light uptake for the practice of photosynthesis, providing better growth and accumulation of dry mass by plants [15]. It is clear that both the excess and the lack of water caused a decrease in the accumulation of DMAP, DMRS and TDM by plants.

The excess water creates anaerobic conditions in the roots, reducing the transport of electrons, damaging the photosynthetic apparatus and the

accumulation of carbohydrates by the roots, limiting the growth of the plants, so under these conditions there is a decrease in the total dry mass production of the plant [20,21,22,23]. Like the excess, the water deficit caused by the lack of water generates inhibition of the photosynthesis by the plants inducing the accumulation of abscisic acid (ABA) promoting the closure of the stomata and limiting the gas exchanges reducing the production of photosimilates and consequently the total dry mass of the plant [17].

The Dickson quality index (DQI) presented quadratic adjustment, with a higher index of 0.29 in the irrigation depth of 8.15 mm d⁻¹ and R² of 0.871 (Fig. 8D). Posse et al. [24], found that the 8 mm d⁻¹ depth is the most adequate for the formation of yellow passion fruit seedlings. Posse et al. [25], also observed that, no matter the climatic conditions, the 8 mm daily depth was the one that provided the best formation of cacao seedlings TSH1188 genotype, with quality and economic use of water and energy.

The DQI is an indicative of quality of seedlings considering in its formula growth characteristics of all parts of the plant. Hunt [26], establishes a minimum DQI value of 0.20 to obtain a quality seedling. Considered also by Johnson and Cline [27], as a promising morphological measure, the DQI reflects the quality of the seedlings considering in its calculation the robustness (TDM) and the balance of the phytomass distribution. The Dickson quality index proved to be a good indicator of quality for the papaya 'Golden THB' seedlings.




Fig. 7. Plant height (A) and stem diameter (B) of papaya 'Golden THB' seedlings submitted to different irrigation depths




Fig. 8. Dry mass of aerial part (A), dry mass of the root system (B), total dry mass (C) and Dickson quality index (D) of papaya 'Golden THB' seedlings submitted to different irrigation depths

In addition, low-DQI seedlings reflect higher mortality rates when they are taken to the field, because they present lower lignification, which would imply a lower survival capacity, thus, seedlings with higher indexes are more likely to survive [28,29].

4. CONCLUSION

The irrigation depths interfered in the growth and quality of papaya 'Golden THB' seedlings during their production period.

For production of papaya 'Golden THB' seedlings with high quality, the indicated irrigation depth is 8.15 mm d⁻¹ as it presented the highest Dickson quality index as well as good responses in the morphological characteristics studied: length of the root system, leaf area, plant height, stem diameter, dry mass of the aerial part, dry mass of the root system and total dry mass.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Serrano LAL, Cattaneo LF. Papaya culture in Brazil. *Revista Brasileira de Fruticultura*, Jaboticabal. 2010;32(3):657-959.
2. Weckner FC, Campos MCC, Nascimento EP, Mantovanelli BC, Nascimento MF. Evaluation of papaya seedlings under the effect of applying different compositions of biofertilizers. *Revista da Universidade Vale do Rio Verde*. 2016;14(1):700-706.
3. Tatagiba SD, Xavier TMT, Torres H, Pezzopane JEM, Cecilio RA, Zanetti SS. Determination of maximum capacity of water retention in substrate for production plants in *Eucalyptus* nursery. *Revista Floresta*. 2015;45(4):745-754.
4. Gruber YBG. Optimization of irrigation depth in the production of seedlings clones of eucalypts (*Eucalyptus urophylla* x *Eucalyptus grandis* and *Eucalyptus urophylla* var. *Platyphylla*). Master Dissertation. Luiz de Queiroz College of Agriculture: University of São Paulo. Piracicaba. 2006;144.

5. Dassie LA, Aleman CC, Moreira ACM, Mignacca FA, Zanfolin PRL, Carvalho PR. Irrigated production of pepper finger seedlings (*Capsicum baccatum*). *Colloquium Agrariae*. 2017;13(3):128-131.
6. Lopes JLW, Guerrini IA, Saad JCC. Effects of irrigation depths on *Eucalyptus grandis* w. (hill ex. maiden) Seedlings in coconut fiber substrate. *Irriga, Botucatu*. 2005; 10(2):123-134.
7. Delgado LGM, Silva RBG, Silva MR. Morphological quality of *Inga vera* seedlings under different water managements. *Irriga, Botucatu*. 2017;22(3):420-429.
8. Köppen W. Das geographische system der klimate. In: Köppen GW, Geiger MR. (eds.), *Handbuch der Klimatologie*, Berlin. 1936;44.
Available:http://koeppen-geiger.vu-wien.ac.at/pdf/Koppen_1936.pdf
9. Allen RG, Pereira LS, Raes D, Smith M. *Crop evapotranspiration: Guidelines for computing crop water requirements. (Irrigation and Drainage Paper, 56)* Rome: FAO. 1998;1-300.
10. Dickson A, Leaf AL, Hosner JF. Quality appraisal of white spruce and white pine seedling stock in nurseries. *Forestry Chronicle*. 1960;36:10-13.
11. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria; 2017.
Available:<https://www.R-project.org/>
12. Trindade AV. Papaya, Production: Technical aspects. *Embrapa Mandioca e Fruticultura*, Brasília, Comunicado para transferência de tecnologia. 2000;77.
13. Severino LS, Cardoso GD, Vale LS, Santos JW. A method for castor bean leaf area estimation. *Rev. Bras. Ol. Fibros. Campina Grande*. 2004;8(1):753-762.
14. Sales RA, Sales RA, Nascimento TA, Silva TA, Berilli SS, Santos RA. Influence of different sources of organic matter in the spread of *Schinus terebinthifolius* Raddi. *Revista Scientia Agraria, Curitiba*. 2017;18(4):99-106.
15. Melo AS, Costa CX, Brito MEB, Viégas PRA, Silva Júnior CD. Production of papaya seedlings in different substrates and phosphorus levels. *Brazilian Journal of Agricultural Sciences, Recife*. 2007;2(4): 257-261.
16. Diniz GL, Sales GN, Sousa VFO, Andrade FHA, Silva SS, Nobre RG. Production of papaya seedlings under water salinity irrigation and phosphate fertilization. *Journal of Agricultural Sciences, Lisbon*. 2018;41(1):218-228.
17. Taiz L, Zeiguer E, Moller IM, Murphy A. *Physiology and plant development*. 6. Ed. Porto Alegre: Artmed. 2017;858.
18. Larcher W. *Plant ecophysiology*. São Carlos: Rima; 2000.
19. Carneiro JGA. *Production and quality control of seedlings*. Curitiba: UFPR/FUPEF. 1995;451.
20. Martinazzo EG, Perboni AT, Oliveira PV, Bianchi VJ, Bacarin MA. Photosynthetic activity in Japanese plum under water deficit and flooding. *Ciência Rural, Santa Maria*. 2013;43(1):35-41.
21. Mazzuchell EL, Souza GM, Pacheco AC. Hardening of eucalyptus seedlings via salicylic acid application. *Pesquisa Farming Tropical, Goiânia*. 2014;44(4): 443-450.
22. Silveira HRO, Souza KRD, Alves JD, Santos MO, Andrade CA, Bomfim SC. Gas exchange and carbohydrate partitioning in coffee seedlings under waterlogging. *Ciênc. Agrotec. Lavras*. 2015;39(2):138-146.
23. Queiroz TB, Rocha SMG, Fonseca FSA, Alvarenga ICA, Martins ER. Effects of water stress on production of eucalypt seedlings. *Irriga, Botucatu*. 2017;22(4): 659-674.
24. Posse RP, Valani F, Gonçalves MAS, Oliveira EC, Louzada JM, Quartezani WZ, Leite MCTL. Growth and quality of yellow passion fruit seedlings produced under different irrigation depths. *Journal of Experimental Agriculture International*. 2018;22(4):1-11.
Available:<http://dx.doi.org/10.9734/jeai/2018/40948>
25. Posse RP, Valani F, Souza CAS, Silveira SS, Silva SMF, Partelli RL, Trindade IM, Costa GS. Growth and quality of genotype TSH1188 cacao tree seedlings produced under different seasons and irrigation depths. *Journal of Experimental Agriculture International*. 2018;28(2):1-17.
Available:<http://dx.doi.org/10.9734/jeai/2018/44718>
26. Hunt GA. Effect of styroblock design and cooper treatment on morphology of conifer seedlings. In: Rose R, Campbell SJ, Landis TD, eds. *Proceedings, Western Forest Nursery Association; 1990 August 13-17; Roseburg, OR. General Technical*

- Report RM-200. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 1990;218-222. Available:<https://rngr.net/publications/proceedings/1990/hunt.pdf/> (Accessed 10 Oct. 2017)
27. Johnson JD, Cline ML. Seedling quality of Southern pines. In: Duryea ML, Dougherty PM (eds.). Forest regeneration manual. Klumer Academic, Netherlands. 1991;143-162. Available:http://dx.doi.org/10.1007/978-94-011-3800-0_8
28. Gomes JM, Couto L, Leite HG, Xavier A, Garcia SLR. Growth of *Eucalyptus grandis* seedlings produced in different sized tubes and N-P-K fertilization. Revista Árvore, Viçosa. 2003;27(2):113-127.
29. Binotto AF, Lúcio ADC, Lopes SJ. Correlations between growth variables and the Dickson quality index in forest seedlings. Revista Cerne, Lavras. 2010;16(4):457-464.

© 2019 Posse et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46286>